



## 1. Introduction

The Water Authorities of North Spain (Confederación Hidrográfica del Cantábrico y del Miño-Sil) are responsible for the management of the Northern Spanish Rivers (Figure 1). During the WFD implementation process and application of the technical annexes the CHC-MS have developed new ecological classification systems following WFD requirements for invertebrates and diatoms (Pardo et al., 2007). There have been also several projects and phases of data collection to fulfil WFD demands in relation with hydromorphological and physico-chemical quality conditions accompanying the high and good status classes, in order to integrate the final ecological classification of water bodies.

In this presentation we explain the approach used in the characterisation of the hydromorphological conditions accompanying the high status based on the invertebrate component. To do it, we have used hydromorphological variables gathered at different scales, from basin to reach and to habitat, to predict the ecological class derived from the invertebrate component. In the future the aim is for integrating the evaluation coming from the fish ecological classification.

## 2. Study area



Figure 1. River typology and monitoring networks of the North of Spain.

## 3. Methodological approach

### Proposed criteria for assessing hydromorphological risk




Table 1. Approximation used in the integration of the hydromorphological criteria to the high status.

## 4. Data bases

Scale	Variables in model	Explanation of variables
Basin	Channelisation rivers_basin	% of meters channelised per basin
	Dams_basin	% of dams higher than 3 m per basin
	Number of dams	Number of dams higher than 3 m upstream
	Weirs_basin	% of dams lower than 3 m per basin
Basin	Industrial & commercial areas	CORINE land cover 1.2.1.
	Road, rail networks & associates land	CORINE land cover 1.2.2.
	Mining production areas	CORINE land cover 1.3.1.
	Rubbish dumps areas	CORINE land cover 1.3.2.
	Construction areas	CORINE land cover 1.3.3.
	Habitat	IHF
Reach	Stream type	River typology following WFD system B based on invertebrates

Table 2. Variables used in the hydromorphological model

## 4. Hydromorphological model

Relationships between hydromorphological variables and ecological classes were established with logistic regressions, allowing for the extraction of variables that significantly contributed to explain the variation between samples from the ecological classes. The dependent variable is the ecological class and the independent variables are the hydromorphological variables. The model was performed with the programme SPSS v. 14.

The model used initially a database of 580 samples for whose no point pressures (organic) were present, being hydromorphological alteration the dominant pressure. In a second run, we also included other high status sites which were not considered previously (669 samples). The model performed with the 669 samples provided a significant  $R^2=0.68$ , and the global percentage of the model for the 5 ecological classes was of 63.5%. The prediction of classes from Moderate to bad was very low. Mainly due to the fact that sites in the worse classes were affected for more than one dominant pressure, and hydromorphological variables could not account for the total variation in the samples within these classes. The model predictions were very good for the high (89% of correct cross validation) and good classes (61% of correct cross validation) (Table 3).

Observed	Pronosticated					Correct percentage
	High	Good	Moderate	Deficient	Bad	
High	201	24	1	0	0	88.9%
Good	72	140	13	4	0	61.1%
Moderate	12	55	38	16	0	31.4%
Deficient	3	12	12	44	2	60.3%
Bad	2	2	3	11	2	10.0%
Global percentage	43.3%	34.8%	10.0%	11.2%	6%	63.5%

Table 3. Classification with the observed and predicted cases in each ecological class provided by the hydromorphological model, and the percentage of correct predictions for each class and for model.




Fig 1. Relationships between the EQR provided by the invertebrate classification and some of the hydromorphological indicators. The reference line in green represents the boundary between Good and moderate status.

		Exclusion thresholds for High ES
Individual variables	Channelisation rivers_basin	5%
	Dams_basin	0,20%
	Weirs_basin	0%
	Industrial & commercial areas	0,10%
	Road, rail networks & associates land	0,25%
	Mining production areas	2%
	Rubbish dumps areas	0,00%
	Construction areas	0,50%
IHF	59	

Table 4. Provisional exclusion thresholds for the high status extracted for the individual variables from the Hydrological model

## 6. Conclusions

The Hydromorphological model allowed the inclusion of hydromorphological elements of importance for supporting the biological elements, and to check the High status provided by invertebrates, in the integration of the final ecological classification. The presented methodological approach is going to be improved when the fish classification will be finalised, and other variables of importance for the migratory fishes will be included (i.e. downstream river continuity).

The high ecological status can be confirmed by the variables of the model. A small number of sites in High status passed to be in good status according the model prediction. Other sites in good status were considered by the model to be in High status according the hydromorphological elements. This result highlights the potential for the model to identify sites in high HM conditions, indicating that the programme of measures and restoration at the reach-local scale can have a great potential to achieve high status. Also the model results will help in the improvement of the boundaries used up to now for the High/good classes.

## 7. Other projects

- 1.- DELINEATION OF PROTECTED RIPARIAN ZONES IN THE CANTABRIC & MIÑO-SIL WATER DISTRICTS. The Cantabric & Miño-Sil water districts have signed a series of general protocols of cooperation with various councils within its territorial area to establish riparian protection zones with the objective of avoiding future human use of flooding areas linked with river dynamics. Up to know a total of 35 councils are participating.
- 2.- GEOMORPHOLOGIC EVOLUTION OF FLUVIAL REACHES. The rivers have been mapped, together with the existing roads and all lands influenced by the functioning of the rivers. The project studies the natural or artificial succession with historical aerial photograph and geomorphological.
- 3.- Regional projects generating obstacles inventories to assess the impacts of barriers to fish migration.
- 4.- Two phases of IMPRESS data collection to identify significant morphological alterations to water bodies.